

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

CODE: 8220

NAME: Siliceous rocky slopes with chasmophytic vegetation

## 1. National Level

### 1.1 Maps

1.1.1 Distribution Map	Yes
1.1.2 Distribution Method	Estimate based on partial data with some extrapolation and/or modelling (2)
1.1.3 Year or period	2006-2012
1.1.4 Additional map	No
1.1.5 Range Map	Yes

## 2. Biogeographical Or Marine Level

### 2.1 Biogeographical Region

### 2.2 Published

#### Mediterranean (MED)

Dimopoulos P., Xystrakis F. and Tsiripidis I. 2014. Deliverable A1. Final Catalogue of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 54.

Dimopoulos P., Fotiadis G., Tsiripidis I., Panitsa M. and Karadimou E. 2014. Deliverable A2. Report and Literature Database on Habitat Types of Greece – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 210.

Tsiripidis I., Xystrakis F., Kasampalis D., Mastrogianni A., Strid A. and Dimopoulos P., 2014. Deliverable A4. Potential Distribution Maps of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, Athens, pages 176.

Dimopoulos P., Tsiripidis I., Xystrakis F., Panitsa M., Fotiadis G., Kallimanis A.S. and Kazoglou I. 2014. Deliverable A6. Explanatory Implementation Manual for the Conservation Degree Assessment of Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 35. (with Annexes: I. Habitat types protocols, pages 600; II. Explanatory notes on the habitat types protocols selection, pages 4; III. Correspondence of Habitat types protocols with the clusters of vegetation relevés (excel file).

Dimopoulos P., Tsiripidis I., Xystrakis F., Kallimanis A.S and Panitsa M. 2014. Deliverable A7. Preliminary Analysis of the Field Data for the Habitat Types – 1st edition. Ministry of Environment, Energy and Climate Change, OIKOM Ltd - E. Alexandropoulou - A. Glavas, Athens, pages 16.

Βραχνάκης Μ., Φωτιάδης Γ. & Καζόγλου Ι. 2011. Τύποι Οικοτόπων Εθνικού Πάρκου Πρεσπών – Αναγνώριση-Καταγραφή 2011. Εταιρία Προστασίας Πρεσπών, σελ. 101.

Dimopoulos P., Sýkora K.V., Mucina L. & Georgiadis T. 1997. The high-rank syntaxa of the rock-cliff and scree vegetation of the mainland Greece and Crete. *Folia Geobotanica* 32 (3): 313-334.

Ελευθεριάδου Ε., Τσιριπίδης Ι., Θεοδωρόπουλος Κ. & Ξυστράκης Φ. 2007. Τύποι οικοτόπων της περιοχής "Ροδόπη (Σημύδα)" του Δικτύου "Φύση 2000". Πρακτικά 13ου Πανελληνίου Δασολογικού Συνεδρίου της Ελληνικής Δασολογικής Εταιρείας, Χλόη Καστοριάς, 7-10 Οκτωβρίου 2007 (τόμος Ι): 91-99.

Θεοδωρόπουλος Κ., Ελευθεριάδου Ε., Τσιριπίδης Ι. & Αθανασιάδης Ν. 2001. Βραχύφιλες και λιβαδικές φυτοκοινωνίες του Παρθένου Δάσους Φρακτού του Νομού Δράμας (Α. Μακεδονία, Ελλάδα). Πρακτικά 9ου Πανελληνίου Δασολογικού Συνεδρίου, Κοζάνη, 17-20 Οκτωβρίου 2000: 661-674.

Θεοχαρόπουλος Μ., Δημητράλλος Γ., Χοχλιούρος Σ., Μαρούλης Γ. & Γεωργιάδης

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Θ. 2002. Συμβολή στην μελέτη των ασβεστολιθικών χασμοφυτικών φυτοκοινοτήτων των τάξεων *Onosmetalia frutescentis* Quezel 1964 και *Potentilletalia speciosae* Quezel 1964 στην ηπειρωτική Ελλάδα. Πρακτικά 9ου Πανελληνίου Επισημονικού Συνεδρίου της Ελληνικής Βοτανικής Εταιρίας, Αργοστόλι Κεφαλονιάς, 9-12 Μαΐου 2002: 129-135.

Quézel P. 1969. La végétation du massif du Bela-Voda (Macédoine nord-occidentale). *Biol. Gallo-Hellen.* 2(2): 93-112.

## 2.3 Range of the habitat type in the biogeographical region or marine region

2.3.1 Surface area - Range (km <sup>2</sup> )	43,43
2.3.2 Range method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.3.3 Short-term trend period	2001-2012
2.3.4 Short-term trend direction	stable (0)
2.3.5 Short-term trend magnitude	min max
2.3.6 Long-term trend period	
2.3.7 Long-term trend direction	N/A
2.3.8 Long-term trend magnitude	min max
2.3.9 Favourable reference range	area (km <sup>2</sup> ) operator approximately equal to (≈) unknown No method
2.3.10 Reason for change	Improved knowledge/more accurate data Use of different method

## 2.4 Area covered by Habitat

2.4.1 Surface area (km <sup>2</sup> )	43,43
2.4.2 Year or period	2000-2012
2.4.3 Method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.4 Short-term trend period	2001-2012
2.4.5 Short-term trend direction	stable (0)
2.4.6 Short-term trend magnitude	min max
2.4.7 Short term trend method used	Estimate based on partial data with some extrapolation and/or modelling (2)
2.4.8 Long-term trend period	
2.4.9 Long-term trend direction	N/A
2.4.10 Long-term trend magnitude	min max
2.4.11 Long term trend method used	N/A
2.4.12 Favourable reference area	area (km) operator approximately equal to (≈) unknown No method
2.4.13 Reason for change	Improved knowledge/more accurate data Use of different method

## 2.5 Main Pressures

Pressure	ranking	pollution qualifier(s)
grazing (A04)	low importance (L)	N/A
Biocenotic evolution, succession (K02)	low importance (L)	N/A

2.5.1 Method used – pressures mainly based on expert judgement and other data (2)

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

## 2.6 Main Threats

Threat	ranking	pollution qualifier(s)
grazing (A04)	low importance (L)	N/A
Biocenotic evolution, succession (K02)	low importance (L)	N/A

2.6.1 Method used – threats expert opinion (1)

## 2.7 Complementary Information

### 2.7.1 Species

Allium flavum

Anthemis orbelica

Asplenium septentrionale

Campanula rotundifolia

Campanula sartorii

Centaurea alba

Centaurea deustiformis

Centaurea ptarmicoides (syn: Centaurea ptarmicifolia)

Euphrasia liburnica

Festuca hirtovaginata

Festuca paniculata

Genista lydia

Globularia cordifolia

Hypericum cuisinii

Minuartia hirsuta

Minuartia recurva

Plantago holosteum

Poa bulbosa

Poa thessala

Polygonum icaricum

Potentilla argentea

Rorippa thracica

Saxifraga paniculata

Scleranthus perennis

Sedum alpestre

Sedum annuum

Sedum sediforme

Sedum stefco

Sempervivum heuffelii

Sempervivum marmoreum

Atocion lerchenfeldianum (syn: Silene lerchenfeldiana)

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

Silene pusilla

Campanula cretica (syn: Symphyandra cretica)

Thymus praecox

Thymus thracicus

Viola doerfleri

## 2.7.2 Species method used

Typical species were determined on the basis of a vegetation database, comprised of about 22000 sampling plots. First, a list of possible typical species was determined per habitat type, selecting the ones presenting a high fidelity value to the habitat types according the algorithm of Tsiripidis et al. (2009) and the phi coefficient value (Chytrý et al. 2002). Then typical species per habitat type were selected from the above-mentioned lists by expert judgment and using as criteria species niche breadth, their ability to comprise indicators of habitat types' conservation status and their function as keystone species. The nomenclature of the typical species follows Dimopoulos et al.

(2013).References Chytrý, M., Tichý, L., Holt, J. & Botta-Duká t, J. 2002. Determination of diagnostic species with statistical fidelity measures. Journal of Vegetation Science 13: 79–90. Dimopoulos, P., Raus, Th., Bergmeier, E., Constantinidis, Th., Iatrou, G., Kokkini, S., Strid, A. & Tzanoudakis, D. 2013: Vascular plants of Greece: an annotated checklist. – Berlin: Botanischer Garten und Botanisches Museum Berlin-Dahlem, Freie Universität Berlin; Athens: Hellenic Botanical Society. Englera 31: 1-367. Tsiripidis, I., Bergmeier, E., Fotiadis, G. & Dimopoulos, P. 2009. A new algorithm for the determination of differential taxa. Journal of Vegetation Science 20: 233-240.

## 2.7.3 Justification of % - thresholds for trends

## 2.7.4 Structure and functions - methods used

Complete survey/Complete survey or a statistically robust estimate (3)

## 2.7.5 Other relevant information

## 2.8 Conclusions (assessment of conservation status at end of reporting period)

2.8.1 Range assessment Favourable (FV)  
qualifiers N/A

2.8.2 Area assessment Favourable (FV)  
qualifiers N/A

2.8.3 Specific structures and functions (incl Species) assessment Favourable (FV)  
qualifiers N/A

2.8.4 Future prospects assessment Favourable (FV)  
qualifiers N/A

2.8.5 Overall assessment of Conservation Status Favourable (FV)

2.8.5 Overall trend in Conservation Status N/A

## 3. Natura 2000 coverage conservation measures - Annex I habitat types on biogeographical level

### 3.1 Area covered by habitat

3.1.1 Surface area (km<sup>2</sup>) min 41,4 max 41,4

# Report on the main results of the surveillance under article 17 for annex I habitat types (Annex D)

3.1.2 Method used

Complete survey/Complete survey or a statistically robust estimate (3)

3.1.3. Trend of surface area

stable (0)

## 3.2 Conservation Measures

3.2.1 Measure

3.2.2 Type

3.2.3 Ranking

3.2.4 Location

3.2.5 Broad Evaluation

Establish protected areas/sites (6.1)

Legal  
Administrative  
One-off

high importance  
(H)

Inside

Maintain  
Long term